

آشنایی با نرم افزار

End Note


بخش ۴

نسخه X6


حمیده حاجی امیری، کارشناس کتابخانه مرکزی دانشگاه

اردیبهشت ۱۳۹۵


وارد کردن منابع به متن مقاله در محیط Word ابزارها و تنظیمات آن



AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period. MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3. Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35?36 minutes. Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.




AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period. MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3. Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35-36 minutes. Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.





AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period. MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3. Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35?36 minutes. Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.





AIM: The need to manage children in paediatric dentistry is high. ketamine (5 mg/kg) and diazepam dental procedures at a paediatric children who were scheduled received ketamine 5 mg/kg b for use as conscious sedation and duration of surgical procedure. Twenty five patients participated action was 10.5 +/- 7.2 minutes cases (12.0%) developed high hallucination. CONCLUSION: The duration of effectiveness sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.



materials and techniques of low dose en undergoing paediatric AND METHODS: All in the study. All children ight in a single oral dose s 2 or 3. Time of onset were recorded. RESULTS: The mean time of onset of 35?36 minutes. Three as a case (4.0%) of was considered effective. for use for conscious

Reference | Preview |  Ashley Paul-2012-Preoperative analges.pdf | 


Reference Type: Electronic Article 

 Hide Empty Fields

Rating
* * * * *

Author
Ashley Paul, F.
Parekh, Susan
Moles David, R.
Anand, Prabhleen
Rehbehani Amal

اسلايدهاي کارگاه هاي آموزشي کتابخانه مرکزي ع.پ. مشهد



AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period. MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3. Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35?36 minutes. Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.

AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period(1-7). MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3. Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35.36 minutes. Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 30 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatient undergoing minor oral surgical procedures.

1. Ashley Paul F, Parekh S, Moles David R, Anand P, Behbehani A. Preoperative analgesics for additional pain relief in children and adolescents having dental treatment. Cochrane Database of Systematic Reviews [Internet]. 2012; (9). Available from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008392.pub2/abstract>
<http://onlinelibrary.wiley.com/store/10.1002/14651858.CD008392.pub2/asset/CD008392.pdf?v=1&t=hyqqh00j&s=1b01f673eb975b6e5c630b01dd151b3729a0392b>.
2. Askelson NM, Chi DL, Momany E, Kuthy R, Ortiz C, Hanson JD, et al. Encouraging early preventive dental visits for preschool-aged children enrolled in Medicaid: Using the Extended Parallel Process Model to conduct formative research. Journal of Public Health Dentistry. 2014;74(1):64-70.
3. Baelum V. Dentistry and population approaches for preventing dental diseases. Journal of Dentistry. 2011 12//;39, Supplement 2(0):S9-S19.
4. Bridgman C. Commissioning explained: early thoughts on taking dentistry forward with a single commissioner. Faculty of Dental Surgery. 2015;4(3):138-41.
5. Cuadros-Fernandez C, Armengol-Olivares A, Guinet-Limeno E, Lorenzo-Rodriguez AL. Oral



ویرایش رفرنس ها در متن مقاله از طریق Edit & Manage citations


Edit & Manage Citation(s)

Edit formatted single or multiple citations by omitting author and/or year information, or adding prefix or suffix text (such as page numbers).

EndNote Cite While You Write
 Press F1 for add-in help.


AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period(1-7). MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3(8-12). Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35?36 minutes. Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.

1. Ashley Paul F, Parekh S, Moles David R, Anand P, Behbehani A. Preoperative analgesics for additional pain relief in children and adolescents having dental treatment. Cochrane Database of Systematic Reviews [Internet]. 2012; (9). Available from:
<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008392.pub2/abstract>
<http://onlinelibrary.wiley.com/store/10.1002/14651858.CD008392.pub2/asset/CD008392.pdf?v=1&t=hvqgh00j&s=1b01f673eb975b6e5c630b01dd151b3729a0392b>.
2. Askelson NM, Chi DL, Momany E, Kuthy R, Ortiz C, Hanson JD, et al. Encouraging early preventive dental visits for preschool-aged children enrolled in Medicaid: Using the Extended Parallel Process





مرحله نهایی: روزآمدسازی رفرنس ها در متن از طریق Update citation & bibliography



Update Citations and Bibliography

Format (or reformat) your document according to the rules of the selected Bibliographic Output style. The EndNote formatting process replaces the temporary citations in your paper with formatted citations, and builds a bibliography at the end of the document.

 **EndNote Cite While You Write**
Press F1 for add-in help.

AIM in ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period(6-12). MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3. Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35±36 minutes. Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.

1. GarciaLopes R, MendesPinto M, deGodoy CH, Jansiski Motta L, CarvalhoBortoletto C, Olivan S, et al. Aesthetic and functional rehabilitation of child using mock-up combined with stratified technique. European journal of paediatric dentistry : official journal of European Academy of Paediatric Dentistry. 2014 Jul;15(2 suppl):234-6. PubMed PMID: 25101511. Epub 2014/08/08. Eng.
2. Klingberg G, Andersson-Wenckert I, Grindejford M, Lundin S-A, Ridell K, Tsilingaridis G, et al. Specialist paediatric dentistry in Sweden 2008 - a 25-year perspective. International Journal of Paediatric Dentistry. 2010;20(5):313-21.
3. Kühnisch J, Heitmüller D, Thiering E, Brockow I, Hoffmann U, Neumann C, et al. Proportion and

ارسال فایل غیر قابل ویرایش به مجله مورد نظر از طریق گزینه

❖ Convert to plain text

Document1 - Microsoft Word

FileHomeInsertPage LayoutReferencesMailingsReviewViewEndNote X6

EN Go to EndNote

Insert Citation

Edit & Manage Citation(s)

Edit Library Reference(s)

Citations

Style: Vancouver

Update Citations and Bibliography

Convert Citations and Bibliography

Convert to Unformatted Citations

Convert to Plain Text

Convert Reference M

Convert Word Citati

Export to EndNote

Preferences

EndNote Help

Convert to Plain Text

Save a copy of your document without formatted Cite While You Write field codes. Formatted citations and the bibliography are saved as text.

EndNote Cite While You Write

Press F1 for add-in help.


AIM: The need to manage expensive conscious sedation materials and techniques in paediatric dentistry is h and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period(1-7). MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3. Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35?36 minutes(8-12). Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.

1. Ashley Paul F, Parekh S, Moles David R, Anand P, Behbehani A. Preoperative analgesics for additional pain relief in children and adolescents having dental treatment. Cochrane Database of Systematic Reviews [Internet]. 2012; (9). Available from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008392.pub2/abstract>


<http://onlinelibrary.wiley.com/store/10.1002/14651858.CD008392.pub2/asset/CD008392.pdf?v=1&t=hqqh00j&s=1b01f673eb975b6e5c630b01dd151b3729a0392b>.

2. Askelson NM, Chi PL, Momeni E, Kutlay R, Ortiz C, Hanson JD, et al. Encouraging early preventive dental visits for preschool-aged children enrolled in Medicaid: Using the Extended Parallel Process

AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures. children who were received ketamine for use as conscious and duration of sedation. Twenty five patients action was 10.5 +/- Three cases (12.0%) developed high temperature in the night or the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.



1. Ashley Paul F, Parekh S, Moles David R, Anand P, Behbehani A. Preoperative analgesics for additional pain relief in children and adolescents having dental treatment. Cochrane Database of Systematic Reviews [Internet]. 2012; (9). Available from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008392.pub2/abstract>
<http://onlinelibrary.wiley.com/store/10.1002/14651858.CD008392.pub2/asset/CD008392.pdf?v=1&t=hyqqh00j&s=1b01f673eb975b6e5c630b01dd151b3729a0392b>.
2. Askelson NM, Chi DL, Momany E, Kuthy R, Ortiz C, Hanson JD, et al. Encouraging early preventive dental visits for preschool-aged children enrolled in Medicaid: Using the Extended Parallel Process



ive conscious sedation materials and techniques
effectiveness of a combination of low dose
sedation in healthy children undergoing paediatric
ear period(1-7). MATERIALS AND METHODS: All
and 2012 were included in the study. All children
am 0.2 mg/kg body weight in a single oral dose
in the Ramsey Score was 2 or 3. Time of onset
ng and after discharge were recorded. RESULTS:
e sedation was 84.0%. The mean time of onset of
ation needed this after 35.36 minutes(8-12).
day of the procedure. There was a case (4.0%) of

hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective.
The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious
sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.

1. Ashley Paul F, Parekh S, Moles David R, Anand P, Behbehani A. Preoperative analgesics for additional pain relief in children and adolescents having dental treatment. Cochrane Database of Systematic Reviews [Internet]. 2012; (9). Available from:

<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008392.pub2/abstract>

<http://onlinelibrary.wiley.com/store/10.1002/14651858.CD008392.pub2/asset/CD008392.pdf?v=1&t=hvqgh00j&s=1b01f673eb975b6e5c630b01dd151b3729a0392b>

2. Askelson NM, Chi DL, Momany E, Kuthy R, Ortiz C, Hanson JD, et al. Encouraging early preventive




AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3-year period(1-7). MATERIALS AND METHODS: All children who were scheduled for conscious sedation between 2009 and 2012 were included in the study. All children received ketamine 5 mg/kg body weight in combination with diazepam 0.2 mg/kg body weight in a single oral dose for use as conscious sedation. Patients were considered sedated when the Ramsey Score was 2 or 3. Time of onset and duration of surgical procedures were recorded. Side effects during and after discharge were recorded. RESULTS: Twenty five patients participated in the study. The effectiveness of the sedation was 84.0%. The mean time of onset of action was 10.5 +/- 7.2 minutes. All cases that needed additional sedation needed this after 35?36 minutes(8-12). Three cases (12.0%) developed high temperature in the night of the day of the procedure. There was a case (4.0%) of hallucination. CONCLUSION: Ketamine and diazepam as medication for conscious sedation was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.


1. Ashley Paul F, Parekh S, Moles David R, Anand P, Behbehani A. Preoperative analgesics for additional pain relief in children and adolescents having dental treatment. Cochrane Database of Systematic Reviews [Internet]. 2012; (9). Available from:
<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008392.pub2/abstract>

<http://onlinelibrary.wiley.com/store/10.1002/14651858.CD008392.pub2/asset/CD008392.pdf?v=1&t=hqqh00j&s=1b01f673eb975b6e5c630b01dd151b3729a0392b>.

2. Askelson NM, Chir DL, Momany ES, Kuthy R, Ortiz C, Hanson DD, et al. Encouraging early preventive dental visits for preschool-aged children enrolled in Medicaid: Using the Extended Parallel Process



AIM: The need to manage children using safe, effective and inexpensive conscious sedation materials and techniques in paediatric dentistry is high. This study evaluated the safety and effectiveness of a combination of low dose ketamine (5 mg/kg) and diazepam (0.2 mg/kg) used for conscious sedation in healthy children undergoing paediatric dental procedures at a paediatric dental outpatient clinic over a 3 year period(1-7). MATERIALS AND METHODS: All children who were scheduled for a procedure received ketamine 5 mg/kg for use as conscious sedation and duration of surgical procedure was recorded. RESULTS: Twenty five patients participated in the study. All children received a single oral dose of ketamine and diazepam for 3. Time of onset of sedation was recorded. RESULTS: The mean time of onset of sedation was 3.6 minutes(8-12). There was a case (4.0%) of hallucination. CONCLUSION: The combination of ketamine and diazepam was considered effective. The duration of effectiveness appears to be 35 minutes. The combination is considered safe for use for conscious sedation in healthy paediatric dental outpatients undergoing minor oral surgical procedures.



1. Ashley Paul F, Parekh S, Moles David R, Anand P, Behbehani A. Preoperative analgesics for additional pain relief in children and adolescents having dental treatment. Cochrane Database of Systematic Reviews [Internet]. 2012; (9). Available from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008392.pub2/abstract>
<http://onlinelibrary.wiley.com/store/10.1002/14651858.CD008392.pub2/asset/CD008392.pdf?v=1&t=hqqh00J&s=1b01f673eb975b6e5c630b01dd151b3729a0392b>
اسلایدهای کارگاه آموزشی آموزش کارگاه آموزشی
2. Askelson NM, Chi DL, Momany E, Kuthy R, Ortiz C, Hanson JD, et al. Encouraging early preventive

چنانچه پرسشی دارید به نشانی زیر ایمیل بزنید:
Cen.lib@mums.ac.ir
کتابخانه مرکزی دانشگاه علوم پزشکی مشهد

اسلایدهای کارگاه های آموزشی کتابخانه مرکزی ع.پ. مشهد